

Europäisches Patentamt European Patent Office Office européen des brevets



(11) EP 0 974 571 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication: 26.01.2000 Bulletin 2000/04

(51) Int Cl.7: C07C 17/25

(21) Application number: 99305781.9

(22) Date of filing: 21.07.1999

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 21.07.1998 US 119560

(71) Applicant: ELF ATOCHEM NORTH AMERICA, INC. Philadelphia, Pennsylvania 19103-3222 (US)

(72) Inventors:

• Elsheikh, Maher Yousef Wayne, Pennsylvania 19087 (US)

 Fellenger, Paul David Narvon, Pennsylvania 17555 (US)

(74) Representative: Stoner, Gerard Patrick et al MEWBURN ELLIS York House 23 Kingsway London WC2B 6HP (GB)

(54) Preparation of 1,1,1,3-tetrafluoropropene(1234ze)

(57) Disclosed is a process for the preparation of cis/trans 1,1,1,3-tetrafluoro-2-propene (1234 ze) which comprises (a) contacting 1,1,1,3,3-pentafluoropropane (245fa) with an alkaline solution, preferably an aqueous

or alcoholic solution of a base such as KOH, NaOH, Ca (OH)₂ or Mg(OH)₂, or with a chromium-based catalyst, such as fluorided Cr₂O₃ or fluorided Cr/Ni/Alf₃, and (b) recovering cis/trans 1,1,1,3-tetrafluoro-2-propene from the reaction mixture.

[0001] This invention relates to preparation of cis/ trans 1,1,1,3-tetrafluoropropene ("1234ze"), a monomer useful for the preparation of various homopolymers and copolymers, particularly to processes for the dehy-1,1,1,3,3-pentafluoropropane drofluorination of ("245fa"), a known blowing agent, to 1234ze using a chromium-based catalyst or a strong base. While the prior art, R. N. Haszeldine, J. Chem. Soc., 1952 (3490), describes the synthesis of cis/trans 1,1,1,3-tetrafluoropropene by fluorination of 1,1,1-trifluoro-2-propyne, this latter feed stock material is not available commercially. [0002] Herein provided is a process for the preparation of 1234ze which comprises

1

- (a) contacting 245fa with an alkaline solution or with a chromium-based catalyst, and
- (b) recovering cis/trans 1,1,1,3-tetrafluoro-2-propene from the resulting reaction mixture.

[0003] It has now been discovered that the cis and trans isomers of 1,1,1,3-tetrafluoro-2-propene (1234ze) can be conveniently prepared by dehydrofluorination of the blowing agent, 245fa, using either a strong base (either an aqueous or alcoholic solution) or a chromiumbased catalyst.

[0004] The catalyzed process is preferably carried out in the gas phase. Use of an oxygen-containing gas such as air is desired to extend the catalyst lifetime, the level of oxygen generally being from about 1 to about 10 volume percent (preferably about 2 to 5%), based on the volume of the organic feed. Temperatures of from about 100°C. to about 600°C. are typically used, preferably from about 300°C, to about 400°C. The pressure can be atmospheric. Contact time (total flow rate per catalyst volume) is typically from about 1 to about 60 seconds, preferably from about 20 to 50 seconds. The catalyst is a chromium-based catalyst such as fluorided chromium oxide, Cr₂O₃, which chromium-based catalyst is either unsupported or supported on a support such as activated carbon, graphite, fluorided graphite or fluorided alumina, the chromium catalyst being used alone or in the presence of a co-catalyst selected from a nickel, cobalt, manganese or zinc salt. Two such preferred chromium catalysts are high surface area chromium oxide and chromiuminickel on fluorided alumina (Cr/Ni/AIF3), preparation of this latter catalyst being taught, for example, in European Patent 486333. The chromium-based catalysts are preferably activated before use, typically by a procedure wherein the catalyst bed is heated to about 370°-380°C. (normally with a continuous flow of nitrogen), after which a mixture of approximately equal volumes of HF and air or nitrogen (preferably nitrogen) are fed over the catalyst bed for about 18 hours.

[0005] The dehydrofluorination can also be accomplished using an alkaline solution of a strong base, such as an aqueous or alcoholic solution of potassium hydroxide (KOH), sodium hydroxide (NaOH), calcium hydroxide (Ca(OH)₂) or magnesium hydroxide (Mg(OH)₂). For the alcoholic solution, a conventional alcohol such as ethanol can be used. The solution typically is from about 0.01 to about 10 molar, preferably 0.1 to 5 molar. The dehydrofluorination is typically conducted at a temperature of from about 20°C, to about 100°C, preferably from about 20°C, to about 50°C.

[0006] The following examples are illustrative.

[0007] Example 1. 52.4 Grams of a high surface area Cr₂O₃ catalyst was activated by first feeding 30 ccm of nitrogen for 2 hours at 370°C. followed by cofeeding 30 ccm of HF and 30 ccm of nitrogen for 18 hours at 370°C. Subsequently, a mixture of 20 ccm of 245fa and 3 ccm of air (equal to about 3 volume % of oxygen, based on the 245fa volume) was fed over the catalyst bed at 400°C for a contact time of 45 seconds. Conversion was 96.2%. Selectivity for the desired (1234ze) product was about 96.3% (about 18.5% cis, about 77.8% trans). Performance of the catalyst was steady for 360 hours.

[0008] Example 2. Example 1 was repeated using Cr/ Ni/AIF₃ catalyst (activated at 370°C, using a cofeed of 30 ccm of nitrogen and 30 ccm of HF for 18 hours) in a series of 3 tests, using the same temperature and air/ 245fa feed ratio, but with the contact time between 26 and 39 seconds. Conversions ranged from 88 to 94.5%. Selectivity for the desired (1234ze) product ranged from 96.2 to 98.5% (17.7 to 20.5% cis, 77 to 80.5% trans).

[0009] Example 3. 10 ccm of 245fa was bubbled through 3000 ml of 2.7 molar KOH solution at room temperature (about 20°C.). Analysis of the gaseous dry product, using gas chromatography on line, showed 26% conversion, with selectivity for the desired (1234ze) product of 97.9% (23.9% cis, 74% trans).

Claims

35

40

45

50

- A process for the preparation of cis/trans 1,1,1,3-tetrafluoro-2-propene which comprises (a) contacting 1,1,1,3,3-pentafluoropropane with an alkaline solution or with a chromium-based catalyst, and (b) recovering cis/trans 1,1,1,3-tetrafluoro-2-propene from the resulting reaction mixture.
- 2. A process for the preparation of cis/trans 1,1,1,3-tetrafluoro-2-propene which comprises (a) contacting 1,1,1,3,3-pentafluoropropane with an aqueous or alcoholic solution of base selected from the group consisting of potassium hydroxide, sodium hydroxide, calcium hydroxide or magnesium hydroxide, and (b) recovering cis/trans 1,1,1,3-tetrafluoro-2-propene from the resulting reaction mixture.
- A process as in Claim 2 wherein step (a) comprises contacting 1,1,1,3,3-pentafluoropropane with an aqueous potassium hydroxide solution.

4. A process for the preparation of cis/trans 1,1,1,3-tetrafluoro-2-propene which comprises (a) contacting 1,1,1,3,3-pentafluoropropane with an oxygen-containing gas in the presence of a fluorided catalyst selected from Cr₂O₃ or Cr/Ni/AIF₃, and (b) recovering cis/trans 1,1,1,3-tetrafluoro-2-propene from the resulting reaction mixture.

 A process as in Claim 4 wherein the catalyst is fluorided Cr₂O₃.

 A process as in Claim 4 wherein the catalyst is fluorided Cr/Ni/AIF₃.

15

10

20

25

30

35

40

45

50



Europäisches Patentamt European Patent Office Office européen des brevets



(11) EP 0 974 571 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3: 12.04.2000 Bulletin 2000/15

(51) Int Cl.7: C07C 17/25

(43) Date of publication A2: 26.01.2000 Bulletin 2000/04

(21) Application number: 99305781.9

(22) Date of filing: 21.07.1999

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE
Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 21.07.1998 US 119560

(71) Applicant: ELF ATOCHEM NORTH AMERICA, INC. Philadelphia, Pennsylvania 19103-3222 (US)

(72) Inventors:

Elsheikh, Maher Yousef
 Wayne, Pennsylvania 19087 (US)

 Fellenger, Paul David Narvon, Pennsylvania 17555 (US)

 (74) Representative: Stoner, Gerard Patrick et al MEWBURN ELLIS York House
 23 Kingsway London WC2B 6HP (GB)

(54) Preparation of 1,1,1,3-tetrafluoropropene(1234ze)

(57) Disclosed is a process for the preparation of cis/trans 1,1,1,3-tetrafluoro-2-propene (1234 ze) which comprises (a) contacting 1,1,1,3,3-pentafluoropropane (245fa) with an alkaline solution, preferably an aqueous

or alcoholic solution of a base such as KOH, NaOH, Ca $(OH)_2$ or $Mg(OH)_2$, or with a chromium-based catalyst, such as fluorided Cr_2O_3 or fluorided $Cr/Ni/Alf_3$, and (b) recovering cis/trans 1,1,1,3-tetrafluoro-2-propene from the reaction mixture.

Printed by Jouve, 75001 PARIS (FR)



EUROPEAN SEARCH REPORT

Application Number EP 99 30 5781

		RED TO BE RELEVANT			
Calegory	Citation of document with ind of relevant passa		Relevant to claim	CLASSIFICATION OF THE APPLICATION (INLCI.7)	
x	OLEFINS. COMMUNICATE HYDROGENATION OF PER BULLETIN OF THE ACAR USSR, DIVISION OF CH 1 January 1960 (1960 1312-1317, XP0005788 ISSN: 0568-5230	RFLUORO OLEFINS" DEMY OF SCIENCES OF THE HEMICAL SCIENCES, D-01-01), pages B79 On (XI) -> (XIII) and	1-3	C07C17/25	
A	US 3 579 595 A (REG 18 May 1971 (1971-0		1-3	·	
A	US 3 499 048 A (REG 3 March 1970 (1970-		1-3		
A	Effects in Eliminat Reaction of 2-Aryle JOURNAL OF ORGANIC	CHEMISTRY, pril 1974 (1974-04-05),	1-3	TECHNICAL FIELDS SEARCHED (Int.CI.7) CO7C	
X	WO 96 05157 A (ICI LLEWELLYN (GB); SHA 22 February 1996 (1 * claim 6; example	PLC ; POWELL RICHARD RRATT ANDREW PAUL (GB)) 996-02-22) 6 *	1,4,5		
X	EP 0 234 002 A (PEN 2 September 1987 (1 * claim 1 *	NWALT CORP) 987-09-02)	1,4,5		
X	WO 96 41679 A (DU P (US); COULSTON GEOR 27 December 1996 (1 * page 5, line 1 - 1-5,7-9 *	line 12; examples	1,4,5		
		-/]		
	The present search report has I	been drawn up for all claims			
	Place of search	Date or completion of the search	<u> </u>	Examiner	
	MUNICH	15 February 2000	Ja	anus, S	
CATEGORY OF CITED DOCUMENTS X: particularly relevant if token alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure		E : earlier patent do: after the filing dat her D : document cited i L : document cited fi	T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document ofted in the application L: document cited for other reasons a: member of the same patent family, corresponding document		



EUROPEAN SEARCH REPORT

Application Number EP 99 30 5781

	Citation of document with in	RED TO BE RELEVANT	Relevant	CLASSIFICATION OF THE	
ategory	of relevant passa		to claim	APPLICATION (Int.CI.7)	
(·	PATENT ABSTRACTS OF vol. 003, no. 152 (0 14 December 1979 (19 & JP 54 130507 A (DA 9 October 1979 (1979) * abstract *	C-067), 979-12-14) AIKIN IND LTD),	1,4,5		
D,A	EP 0 486 333 A (ATO 20 May 1992 (1992-0		1,4,6		
Р,Х	WO 98 33756 A (DU P (US); RAO V N MALLI 6 August 1998 (1998 * claim 9; example	ONT ;NAPPA MARIO JOSEPH KARJUNA (US); SIEVE) -08-06) 1 * 	1,4,5		
		·		TECHNICAL FIELDS SEARCHED (Int.Cl.7)	
		·			
		·			
	·	÷			
ox-					
	The present search report has	been drawn up for all claims			
	Place of search	Date of completion of the search		Ехатипет	
X:pa Y:pa do A:te	MUNICH CATEGORY OF CITED DOCUMENTS articularly relevant if taken alone urbicularly relevant if combined with anci- cument of the same category chnological background on-written disolosure	E : earlier patent de after the filing d. ther D : document oited L : document oited	noiple underlying the invention t document, but published on, or y date ted in the application ed for other reasons		

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 99 30 5781

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

15-02-2000

	ocument arch report		Publication date	1	Patent family member(s)	Publication date
US 3579	595	Α	18-05-1971	NONE		
US 3499	048	Α	03-03-1970	US	3564101 A	16-02-1971
WO 9605	157	A	22-02-1996	EP JP US	0775100 A 10505337 T 5856593 A	28-05-1997 26-05-1998 05-01-1999
EP 0234	002	 A	02-09-1987	JР	62169737 A	25-07-1987
WO 9641	679	A	27-12-1996	EP JP	0831965 A 11507870 T	01-04-1998 13-07-1999
JP 5413	80507	A	09-10-1979	JP JP	1283411 C 60007975 B	27-09-198 28-02-198
EP 0486	3333	A	20-05-1992	FR US AT AU CA DE DK ES FI GR JP JP KR MX NO PT	2669022 A 5731481 A 115532 T 641292 B 8779991 A 2055281 A,0 69105924 T 486333 T 2066394 T 915341 A 3015222 T 2053012 C 4288027 A 7080796 B 9506521 B 9102049 A 174096 C 99489 A	15-05-1992 24-03-1994 15-12-1999 16-09-1999 14-05-1999 26-01-1999 27-07-1999 01-05-1999 14-05-1999 14-05-1999 13-10-1999 13-10-1999 16-06-1999 08-07-1999 30-09-1999
	- 3756	Α	06-08-1998	EP US	0968161 A 5945573 A	05-01-200 31-08-199

O w For more details about this annex : see Official Journal of the European Patent Office, No. 12/82